

Rings of definable scalars of some $sl_3(\mathbb{C})$ -modules

Sonia L'Innocente

*School of Science and Technology, Division of Mathematics
University of Camerino, Camerino, Italy*

sonia.linnocente@unicam.it

This a joint work with Mike Prest [4].

In the paper [1], Herzog investigated the ring of definable scalars of the finite-dimensional representations of the Lie algebra $sl_2(\mathbb{C})$ of the 2×2 traceless matrices over the complex field \mathbb{C} . This is the ring of definable actions on the category of finite-dimensional $sl_2(\mathbb{C})$ -modules that is, the ring to which the action of the universal enveloping algebra, $U = U(sl_2(\mathbb{C}))$ on these modules extends in a definable way. Herzog showed, that this ring, denoted by U' is von Neumann regular and is a universal localisation of U . This work inspired further investigations, on rings of definable scalars of Verma modules [3], on $U_q(sl_2(\mathbb{C}))$ -modules (where q is not a root of unity) [2]. It is natural to ask what happens when $sl_2(\mathbb{C})$ is replaced by other simple Lie algebras, in particular by $sl_3(\mathbb{C})$. We are able to obtain the similar results described by [1] if we restrict to the representations which are contained in, or whose dual is contained in, the natural representation of $sl_3(\mathbb{C})$ on the polynomial ring on three generators.

References

- [1] I. Herzog, *The pseudo-finite dimensional representations of $sl(2, k)$* , *Selecta Mathematica*, 7 (2001), 241-290
- [2] I. Herzog, S. L'Innocente, *The Nonstandard quantum plane*, *Annals of Pure and Applied Logic*, 156 (2008), no. 1, 78-85
- [3] S. L'Innocente, M. Prest, *Rings of definable scalars of Verma modules*, *Journal of Algebra and its Applications*, 6 (2007), no. 5, 779-787
- [4] S. L'Innocente, M. Prest, *Rings of definable scalars of some $sl_3(\mathbb{C})$ -modules*, In preparation